

## New England Biolabs Certificate of Analysis

**Product Name:** *EnGen<sup>®</sup> Lba Cas12a (Cpf1)*  
**Catalog #:** *M0653S*  
**Concentration:** *1 μM*  
**Unit Definition:** *A 20 μl reaction in 1X NEBuffer 2.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM crRNA, and 100 nM EnGen<sup>®</sup> Lba Cas12a (Cpf1) incubated for 15 minutes at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.*  
**Lot #:** *0031803*  
**Assay Date:** *03/2018*  
**Expiration Date:** *03/2020*  
**Storage Temp:** *-20°C*  
**Storage Conditions:** *500 mM NaCl, 20 mM Sodium Acetate, 0.1 mM EDTA, 0.1 mM TCEP-HCl, 50% Glycerol, (pH 6.0 @ 25°C)*  
**Specification Version:** *PS-M0653S v1.0*  
**Effective Date:** *31 Oct 2017*

Assay Name/Specification (minimum release criteria)	Lot #0031803
<b>Endonuclease Activity (Nicking)</b> - A 50 μl reaction in NEBuffer 2.1 containing 1 μg of supercoiled PhiX174 RF I DNA and a minimum of 1 pmol of EnGen <sup>®</sup> Lba Cas12a (Cpf1) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 μl reaction in NEBuffer 2.1 containing 1 μg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 1 pmol of EnGen <sup>®</sup> Lba Cas12a (Cpf1) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 μl reaction in NEBuffer 2.1 containing 1 μg of Lambda DNA and a minimum of 1 pmol of EnGen <sup>®</sup> Lba Cas12a (Cpf1) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>RNase Activity (Extended Digestion)</b> - A 10 μl reaction in NEBuffer 4 containing 40 ng of f-300 RNA transcript and a minimum of 1 pmol of EnGen <sup>®</sup> Lba Cas12a (Cpf1) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	<b>Pass</b>



Authorized by  
Derek Robinson  
31 Oct 2017



Inspected by  
Fei Liu  
09 Apr 2018

