

New England Biolabs Certificate of Analysis

Product Name: OneTaq® DNA Polymerase
Catalog Number: M0480L
Concentration: 5,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that will incorporate 15 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.
Packaging Lot Number: 10184640
Expiration Date: 03/2025
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.5 % Tween® 20 , 0.5 % IGEPAL® CA-630 , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0480S/L/X v2.0


OneTaq® DNA Polymerase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0480LVIAL	OneTaq® DNA Polymerase	10182087	Pass
B9026AVIAL	OneTaq® High GC Enhancer	10179724	Pass
B9023SVIAL	OneTaq® GC Reaction Buffer	10157598	Pass
B9022SVIAL	OneTaq® Standard Reaction Buffer	10157597	Pass

Assay Name/Specification	Lot # 10184640
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 units of OneTaq® DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>PCR Amplification (5.0 kb Lambda DNA) A 25 µl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 0.625 units of OneTaq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.</p>	Pass
<p>PCR Amplification (Buffer Dependent, >65% GC-rich) A 25 µl reaction in OneTaq® GC Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 10 ng Human Genomic DNA with 0.625 units of OneTaq® DNA Polymerase for 30 cycles of PCR amplification results in the buffer-dependent production of the expected 737 bp product.</p>	Pass

Assay Name/Specification	Lot # 10184640
<p>PCR Amplification (Enhancer Dependent, >70% GC-rich) A 25 µl reaction in OneTaq[®] GC Reaction Buffer and 20% OneTaq[®] High GC Enhancer in the presence of 200 µM dNTPs and 0.2 µM primers containing 10 ng Human Genomic DNA with 0.625 units of OneTaq[®] DNA Polymerase for 30 cycles of PCR amplification results in the enhancer-dependent production of the expected 627 bp product.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq[®] DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Lea Antonopoulos
Production Scientist
17 Mar 2023



Michael Tonello
Packaging Quality Control Inspector
21 Mar 2023