

## New England Biolabs Certificate of Analysis

Product Name: 7-deaza-dGTP Catalog #: N0445S/L 5 mM Concentration: Unit Definition: N/A0431508 *Lot #:* 08/2015 Assay Date: 08/2017 Expiration Date: -20°C Storage Temp:

Storage Conditions: Supplied in Ultrapure water as a lithium salt, (pH 7.0)

Specification Version: PS-N0445S/L v1.0
Effective Date: 16 Oct 2015

Assay Name/Specification (minimum release criteria)	Lot #0431508
Endonuclease Activity (Nicking) - A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 μl of 7-deaza-dGTP incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 µl of 7-deaza-dGTP incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (0.5 kb Lambda DNA, 7-deaza) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.	Pass
PCR Amplification (2 kb Lambda DNA, 7-deaza) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 2 kb product.	Pass
PCR Amplification (5 kb Lambda DNA, 7-deaza) - A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dATP, dCTP, dTTP and 7-deaza-dGTP, 0.5 μM primers containing 1 ng Lambda DNA with 5 units of Taq® DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.	Pass
<b>Phosphatase Activity (pNPP)</b> - A 200 μl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM <i>p</i> -Nitrophenyl Phosphate (pNPP) and a minimum of 80 μl 7-deaza-dGTP incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass









240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0431508
<b>Physical Purity (HPLC)</b> - 7-deaza-dGTP is ≥ 95% pure as determined by HPLC analysis.	Pass
RNase Activity (Extended Digestion) - A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of 7-deaza-dGTP is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

Authorized by Melanie Fortier 16 Oct 2015







Inspected by Denisa Gilaj 27 Oct 2015