

## New England Biolabs Certificate of Analysis

**Product Name:** Deoxynucleotide (dNTP) Solution Mix  
**Catalog Number:** N0447S  
**Concentration:** 10 mM  
**Unit Definition:** N/A  
**Lot Number:** 10017137  
**Expiration Date:** 07/2020  
**Storage Temperature:** -20°C  
**Storage Conditions:** Supplied in Ultrapure water as a sodium salt (pH 7.5)  
**Specification Version:** PS-N0447S/L v2.0

Deoxynucleotide (dNTP) Solution Mix Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N0447SVIAL	Deoxynucleotide (dNTP) Solution Mix	10017135	Pass

Assay Name/Specification	Lot # 10017137
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Deoxynucleotide (dNTP) Solution Mix is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	<b>Pass</b>
<b>Phosphatase Activity (pNPP)</b> A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl Deoxynucleotide (dNTP) Solution Mix incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	<b>Pass</b>
<b>Physical Purity (HPLC)</b> Deoxynucleotide (dNTP) Solution Mix is ≥ 99% pure as determined by HPLC analysis.	<b>Pass</b>
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 µl of Deoxynucleotide (dNTP) Solution Mix incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b>	<b>Pass</b>

Assay Name/Specification	Lot # 10017137
<p>A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 µl of Deoxynucleotide (dNTP) Solution Mix incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	
<p><b>PCR Amplification (0.5 kb Lambda, dNTPs)</b> A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM Deoxynucleotide (dNTP) Solution Mix and 0.5 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.</p>	<b>Pass</b>
<p><b>PCR Amplification (2.0 kb Lambda, dNTPs)</b> A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM Deoxynucleotide (dNTP) Solution Mix and 0.5 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.</p>	<b>Pass</b>
<p><b>PCR Amplification (5.0 kb Lambda, dNTPs)</b> A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM Deoxynucleotide (dNTP) Solution Mix and 0.5 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.



Cathy Rezac  
Production Scientist  
06 Aug 2018



Michael Tonello  
Packaging Quality Control Inspector  
07 Aug 2018