

New England Biolabs Certificate of Analysis

Product Name: *BstXI*
Catalog Number: *R0113S*
Concentration: *10,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in NEBuffer r3.1 in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10145804*
Expiration Date: *04/2024*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)*
Specification Version: *PS-R0113S/L v2.0*

BstXI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0113SVIAL	BstXI	10145779	Pass
B6003SVIAL	NEBuffer™ r3.1	10132773	Pass

Assay Name/Specification	Lot # 10145804
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of BstXI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) BstXI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of BstXI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of supercoiled pUC19 DNA and a minimum of 30 units of BstXI incubated for 4 hours at 37°C results in <20%</p>	Pass

Assay Name/Specification	Lot # 10145804
conversion to the nicked form as determined by agarose gel electrophoresis.	
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and a minimum of 30 units of BstXI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Functional Testing (15 minute Digest) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and 1 µl of BstXI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with BstXI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BstXI.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Penghua Zhang
Production Scientist
09 May 2022



Michael Tonello
Packaging Quality Control Inspector
09 May 2022