

## New England Biolabs Certificate of Analysis

**Product Name:** BamHI  
**Catalog Number:** R0136L  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10020618  
**Expiration Date:** 08/2020  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R0136S/L v1.0

BamHI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0136LVIAL	BamHI	10020062	Pass
B7203SVIAL	NEBuffer™ 3.1	10018441	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10018419	Pass


Assay Name/Specification	Lot # 10020618
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 20 units of BamHI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Blue-White Screening (Terminal Integrity)</b> A sample of pUC19 vector linearized with a 10-fold excess of BamHI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of BamHI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of BamHI incubated for	Pass

Assay Name/Specification	Lot # 10020618
<p>4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p> <p><b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with BamHI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with BamHI.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.



Tony Spear-Alfonso  
Production Scientist  
21 Aug 2018



Michael Tonello  
Packaging Quality Control Inspector  
31 Aug 2018