

## New England Biolabs Certificate of Analysis

**Product Name:** MscI  
**Catalog Number:** R0534L  
**Concentration:** 5,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10215315  
**Expiration Date:** 12/2025  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % Triton®X-100, 50 % Glycerol, 200 µg/mL rAlbumin (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0534S/L v4.0


MscI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0534LVIAL	MscI	10215290	Pass
B6004SVIAL	rCutSmart™ Buffer	10209244	Pass

Assay Name/Specification	Lot # 10215315
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 units of MscI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with MscI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with MscI.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 25 units of MscI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> MscI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

Assay Name/Specification	Lot # 10215315
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 5 units of MscI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is <math>\leq 1</math> E. coli genome.</p>	<p><b>Pass</b></p>


This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.




---

YunJie Sun  
Production Scientist  
09 Nov 2023




---

Michael Tonello  
Packaging Quality Control Inspector  
20 Dec 2023