

## New England Biolabs Certificate of Analysis

**Product Name:** SexAI  
**Catalog Number:** R0605S  
**Concentration:** 5,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of pBC4 DNA (dcm-) in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10053525  
**Expiration Date:** 07/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R0605S/L v1.0

SexAI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0605SVIAL	SexAI	10050803	Pass
B7204SVIAL	CutSmart® Buffer	10046081	Pass

Assay Name/Specification	Lot # 10053525
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 15 units of SexAI incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Ligation and Recutting (Terminal Integrity)</b>            After a 10-fold over-digestion of pBC4 dcm- DNA with SexAI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with SexAI.</p>	Pass
<p><b>Non-Specific DNase Activity (16 Hour)</b>            A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBC4 dcm- DNA and a minimum of 15 Units of SexAI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Anthony Francis  
Production Scientist  
25 Jul 2019



Jay Minichiello  
Packaging Quality Control Inspector  
28 Aug 2019