

New England Biolabs Certificate of Analysis

Product Name: *Nt.BstNBI*
Catalog Number: *R0607L*
Concentration: *10,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg T7 DNA in NEBuffer r3.1 in 1 hour at 55°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10159560*
Expiration Date: *08/2024*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)*
Specification Version: *PS-R0607S/L v2.0*

Nt.BstNBI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0607LVIAL	Nt.BstNBI	10159559	Pass
B6003SVIAL	NEBuffer™ r3.1	10146826	Pass

Assay Name/Specification	Lot # 10159560
<p>Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of T7 DNA with Nt.BstNBI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Nt.BstNBI.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Nt.BstNBI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Nt.BstNBI incubated for 4 hours at 55°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Non-Specific DNase Activity (16 hour) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of T7 DNA and a minimum of 10 units of Nt.BstNBI incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE:</p>	Pass

Assay Name/Specification	Lot # 10159560
<p>although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.</p> <p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of Nt.BstNBI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Penghua Zhang
Production Scientist
09 Sep 2022



Michael Tonello
Packaging Quality Control Inspector
09 Sep 2022