

## New England Biolabs Certificate of Analysis

*Product Name:* EcoRI-HF<sup>®</sup>  
*Catalog #:* R3101S/L  
*Concentration:* 20,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
*Lot #:* 0141611  
*Assay Date:* 11/2016  
*Expiration Date:* 11/2018  
*Storage Temp:* -20°C  
*Storage Conditions:* 300 mM NaCl, 10 mM KPO<sub>4</sub>, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % TritonX-100, 200 µg/ml BSA, (pH 7.0 @ 25°C)  
*Specification Version:* PS-R3101S/L v2.0  
*Effective Date:* 15 Jun 2016

| Assay Name/Specification (minimum release criteria)   | Lot #0141611 |
|---|--------------|
| <b>Blue-White Screening (Terminal Integrity)</b> - A sample of pUC19 vector linearized with a 10-fold excess of EcoRI-HF <sup>™</sup> , religated and transformed into an <i>E. coli</i> strain expressing the LacZ beta fragment gene results in <1% white colonies.   | <b>Pass</b>  |
| <b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 Units of EcoRI-HF <sup>™</sup> incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.                  | <b>Pass</b>  |
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 200 units of EcoRI-HF <sup>™</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | <b>Pass</b>  |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 20-fold over-digestion of Lambda DNA with EcoRI-HF <sup>™</sup> , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EcoRI-HF <sup>™</sup> .  | <b>Pass</b>  |
| <b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of EcoRI-HF <sup>™</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.     | <b>Pass</b>  |
| <b>Protein Purity Assay (SDS-PAGE)</b> - EcoRI-HF <sup>™</sup> is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.  | <b>Pass</b>  |



---

## New England Biolabs Certificate of Analysis

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



---

Authorized by  
Derek Robinson  
15 Jun 2016



---

Inspected by  
Stephanie Cornelio  
14 Dec 2017

