

New England Biolabs Product Specification

<i>Product Name:</i>	<i>rCutSmart™ Buffer</i>
<i>Catalog #:</i>	<i>B6004S/V</i>
<i>Concentration:</i>	<i>10X Concentrate</i>
<i>Shelf Life:</i>	<i>36 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Composition (1X):</i>	<i>50 mM Potassium Acetate, 20 mM Tris Acetate, 10 mM Magnesium Acetate, 100 µg/ml rAlbumin, (pH 7.9 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-B6004S/V v1.0</i>
<i>Effective Date:</i>	<i>01 Sep 2021</i>

Assay Name/Specification (minimum release criteria)

Conductivity (buffers/solutions) - The conductivity of 10X rCutSmart™ Buffer is between 40 and 46 mS/cm at 25°C.

Endonuclease Activity (Nicking, Buffer) - A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Functional Testing (Restriction Digest, Buffer) - A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda dam- DNA and 1 unit of ClaI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.

Functional Testing (Restriction Digest, Buffer) - A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 unit of MscI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.

Functional Testing (Restriction Digest, Buffer) - A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 unit of EcoRI-HF® incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.

Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

pH (buffers/solutions) - The pH of 10X rCutSmart™ Buffer is between pH 7.8 and 8.0 at 25°C.

RNase Activity (Buffer) - A 10 µl reaction in 1X rCutSmart™ Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.



New England Biolabs Product Specification

*One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.*



Date 01 Sep 2021

Doreen Duquette
Quality Control Manager

