

New England Biolabs Product Specification

<i>Product Name:</i>	<i>Bsa</i> II
<i>Catalog #:</i>	R0536S/L/V
<i>Concentration:</i>	10,000 units/ml
<i>Unit Definition:</i>	One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 60°C in a total reaction volume of 50 µl.
<i>Shelf Life:</i>	24 months
<i>Storage Temp:</i>	-20°C
<i>Storage Conditions:</i>	100 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.01% Triton X-100, 200 µg/ml BSA
<i>Specification Version:</i>	PS-R0536S/L v1.0
<i>Effective Date:</i>	07/29/2013

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 50 units of BsaII incubated for 4 hours at 60°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of Lambda DNA with BsaII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with BsaII.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 50 Units of BsaII incubated for 16 hours at 60°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 07/29/2013

Derek Robinson
Quality Approver

