

New England Biolabs Product Specification

<i>Product Name:</i>	<i>DraIII-HF[®]</i>
<i>Catalog #:</i>	<i>R3510S/L/V</i>
<i>Concentration:</i>	<i>20,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>20 mM Tris-HCl, 300 mM NaCl, 1 mM Dithiothreitol, 0.1 mM EDTA, 200 µg/ml BSA, 50% Glycerol, pH 7.4 @ 25°C</i>
<i>Specification Version:</i>	<i>PS-R3510S/L v1.0</i>
<i>Effective Date:</i>	<i>04/24/2013</i>

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in CutSmart[™] Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 200 units of DraIII-HF[™] incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of Lambda DNA with DraIII-HF[™], ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DraIII-HF[™].

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart[™] Buffer containing 1 µg of Lambda DNA and a minimum of 20 Units of DraIII-HF[™] incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - DraIII-HF[™] is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 04/24/2013

Derek Robinson
Quality Approver

